

BMJ Open Safety of Tdap vaccine in pregnant women: an observational study

Helen Petousis-Harris,¹ Tony Walls,² Donna Watson,¹ Janine Paynter,¹ Patricia Graham,³ Nikki Turner¹

To cite: Petousis-Harris H, Walls T, Watson D, *et al.* Safety of Tdap vaccine in pregnant women: an observational study. *BMJ Open* 2016;**6**:e010911. doi:10.1136/bmjopen-2015-010911

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2015-010911>).

Received 18 December 2015
Revised 29 February 2016
Accepted 4 March 2016



CrossMark

¹Department of General Practice and Primary Health Care, Immunisation Advisory Centre, The University of Auckland, Auckland, New Zealand

²Department of Paediatrics, University of Otago, Christchurch, New Zealand

³Canterbury District Health Board, Christchurch, New Zealand

Correspondence to

Helen Petousis-Harris; h.petousis-harris@auckland.ac.nz

ABSTRACT

Objectives: Actively recruit and intensively follow pregnant women receiving a dose of acellular pertussis vaccine for 4 weeks after vaccination.

Design and settings: A prospective observational study conducted in 2 New Zealand regions.

Participants: Women in their 28th–38th week of pregnancy, recruited from primary care and antenatal clinics at the time of Tdap administration. Telephone interviews were conducted at 48 h and 4 weeks postvaccination.

Main outcomes measures: Outcomes were injection site reactions, systemic symptoms and serious adverse events (SAEs). Where available, data have been classified and reported according to Brighton Collaboration definitions.

Results: 793 women participated with 27.9% receiving trivalent inactivated influenza vaccine concomitantly. 79% of participants reported mild or moderate pain and 2.6% severe pain. Any swelling was reported by 7.6%, induration by 12.0% (collected from 1 site only, n=326), and erythema by 5.8% of participants. Fever was reported by 17 (2.1%) participants, 14 of these occurred within 24 h. Headache, dizziness, nausea, myalgia or arthralgia was reported by <4% of participants, respectively, and fatigue by 8.4%. During the study period, there were 115 adverse events in 113 participants, most of which were minor. At the end of the reporting period, 31 events were classified as serious (eg, obstetric bleeding, hypertension, infection, tachycardia, preterm labour, exacerbation of pre-existing condition and pre-eclampsia). All had variable onset time from vaccination. There were two perinatal deaths. Clinician assessment of all SAEs found none likely to be vaccine related.

Conclusions: Vaccination with Tdap in pregnant women was well tolerated with no SAE likely to be caused by the vaccine.

Trial registration number: ACTRN12613001045707.

INTRODUCTION

In 2011, the US Advisory Committee on Immunization Practices (ACIP) recommended that acellular pertussis-containing vaccine (Tdap) be given to any person, including pregnant women, likely to be in contact with

Strengths and limitations of this study

- This is the largest number of pregnancy exposures to Tdap that have been actively followed up for safety outcomes.
- Our study allows for close examination of common local and systemic vaccine reactions.
- This is an observational study with no direct comparator groups.
- The study is not large enough to explore serious adverse events.
- The women in our group were older (32 years) than the general New Zealand (NZ) maternal population (29.2 years) and more likely to be of NZ European ethnicity (73.5% vs 49.5%). This is most likely a reflection of the health-seeking behaviour associated with these demographics.

infants under the age of 12 months.¹ In September 2012, the UK recommended providing Tdap vaccine for pregnant women ideally between 28 and 38 weeks of pregnancy.² Vaccine administration in pregnancy offers maternal protection against pertussis and also provides for maternal antibody to be passed to the infant, which has been demonstrated to be protective in their first months of life.^{3 4} There are no theoretical safety concerns with administering subunit vaccines in pregnancy and some vaccines, in particular tetanus, have been used widely in pregnant women.⁵ In October 2012, in response to a pertussis epidemic, the New Zealand (NZ) Ministry of Health began funding Tdap vaccine for all women from 28 to 38 weeks gestation.

While there are now several large studies published,^{6–10} at the time of this study, the USA ACIP committee acknowledged that the safety of Tdap immunisation during pregnancy had not been systematically studied, with the only data available coming from small studies, postmarketing surveillance, and the US Vaccine Adverse Event Reporting System.^{1 11}

Our aim was to intensively monitor the safety of Tdap vaccine in a larger group of pregnant women for a period up to 4 weeks postvaccination.

METHODS

Study design, setting and participants

This was a prospective observational study conducted in two different geographical areas of NZ using the same outcome measures and database. There was some difference in recruitment and data collection methods. The Northern arm of the study was conducted primarily in Auckland and included other North Island centres. Auckland is a culturally diverse metropolitan city of 1.42 million in the North Island. Women being administered Tdap between 28 and 38 weeks gestation were identified by staff from 21 out of 24 participating general practices and the maternity clinics of three district health boards (DHBs) and referrals faxed to the study team from 30 January to 30 June 2014.

The other arm was conducted in Canterbury, a more ethnically homogeneous South Island region of just over half a million. Recruitment for this arm began earlier, from late September 2012 to late June 2014. Participants aged 18–40 years and administered Tdap between 30 and 36 weeks gestation were identified via claims submitted by general practices (within 1 week of vaccination) for reimbursement from the local DHB for each vaccination service delivered.

In both study arms, consent to be contacted by members of the research team was sought at the time of vaccination.

Inclusion criterion was being compliant with routine antenatal care, including at least one ultrasound early in pregnancy. Women who had given birth prior to being contacted by study team members were originally to be excluded from both arms of the study; however, owing to concern that this may lead to overlooked serious adverse events (SAEs; ie, premature birth) early in the study, after the first exclusion, subsequent women in the Northern arm who had birthed prior to contact but met other inclusion criteria were included.

Vaccines

The pertussis-containing vaccine (Tdap) funded for pregnant women in NZ is Boostrix. The Boostrix formulation used in NZ has 0.5 mg aluminium as aluminium hydroxide and aluminium phosphate adjuvant and the US formulation has no more than 0.39 mg aluminium as aluminium hydroxide.

Influenza (TIV) vaccine is given at the start of the winter season and funded for all pregnant women regardless of gestational age. Where this time coincides with the gestation age for delivery of the Tdap, providers are encouraged to deliver both vaccines together.

Data collection and main outcomes

Northern participants leaving the practice/clinic after their vaccination were given a study envelope containing an information sheet, consent form, clear plastic measuring tool (to measure any local reaction) and a 3-day diary card to record any symptoms or events. At the first phone contact 48–72 h postvaccine administration, verbal

consent was obtained and an interview undertaken. The second phone interview was conducted at 4 weeks postadministration (see [figure 1](#)).

For Canterbury participants, a research team member made phone contact with potential participants within 2 weeks of identification, obtained consent and conducted the first interview. A follow-up questionnaire was mailed at 4 weeks postvaccination.

For both groups, consent to follow-up with their Lead Maternity Carer (either a midwife or obstetrician) or general practitioner was sought from all women who reported an SAE or any birth complication. As per national protocols,¹² any adverse events (AEs) were also reported to the Centre for Adverse Reaction Monitoring (CARM), the official regulatory body in charge of receiving all AE reports in NZ. An AE is any untoward medical occurrence temporally associated with administration of the vaccine and does not include common injection site reactions. In NZ, AEs following immunisation reporting is encouraged if the event is serious or unexpected. Anonymised data for any SAEs were also reported to the Marketing Authorisation Holder (the vaccine manufacturer) as part of global drug safety surveillance.

To ensure consistency in our classifications of SAEs, we used an algorithm based on the International Conference on Harmonisation definitions for SAEs.^{13 14} We separated events occurring in the mother during pregnancy ([figure 2](#)) and those triggered by indications of non-reassuring fetal status ([figure 3](#)).

An SAE is defined as any untoward medical occurrence that: results in death; is life-threatening; requires inpatient hospitalisation or causes prolongation of existing hospitalisation; results in persistent or significant disability/incapacity; is a congenital anomaly/birth defect; or requires intervention to prevent permanent impairment or damage.

The term ‘life threatening’ used here refers to an event in which the patient was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe.¹⁵

All SAEs were followed up by a study clinician.


Solicited and unsolicited outcomes

The diary card (Northern arm) allowed recording of injection site reactions, fever and unsolicited events over 3 days (days 0, 1 and 2). Response categories were consistent with Brighton Collaboration definitions and guidelines for events following immunisation.^{16–19}

The 48 h postvaccination telephone interview (Northern arm) asked about nausea, vomiting, diarrhoea, headache, fatigue and contact with healthcare services. Participants were also asked if they had any other concerns after the vaccination. The telephone interview conducted by the Canterbury team sought the same information retrospectively for up to 7 days postvaccination.

The 4-week postvaccination telephone interview (Northern arm) and mailed questionnaire (Canterbury

Figure 1 Participant recruitment and data collection.



arm) asked if any events, including contact with health professionals, had occurred since the last interview. Canterbury participants were phoned if their questionnaire was not returned.

Statistical analysis

The size of our sample was based on the number of births in the study regions (28 000 per annum), estimated uptake of pertussis vaccine in pregnancy and the number of women available for follow-up over the study time frame.

Since the two regions recruited and collected data in different ways, we have not attempted to analyse any differences.

Frequencies, percentages and cross-tabulations for demographic and outcome variables were produced using SAS Enterprise Guide V.6. Results are summarised as counts and percentages for each outcome. Values

within outcome variables (eg, pain, redness, swelling and induration) are based on Brighton Collaboration definitions of outcomes. Other outcomes, such as fever, nausea and myalgia, are summarised as presence or absence (ie, yes or no). Missing values are reported in the tables where relevant.

RESULTS

Twenty-four general practices out of 34 invited (70.6%) and three invited DHB maternity clinics were recruited to the Northern arm. Twenty-one of the practices recruited participants. Practices which declined said they were too busy or had too few patients who met the required criteria. In the Canterbury arm, claim forms for financial reimbursement of delivery of the vaccine for 1212 women were received. Attempts to contact women within 2 weeks of their claim form being received by the study team meant 710 women were




Figure 2 Classification of maternal adverse events.

contacted by phone, of whom 467 consented (65.8%). Participant flow is presented in [figure 4](#).

In total, there were 793 participants across both arms, predominantly of NZ European ethnicity (73.5%). The mean age was 32, with 61.4% of participants aged between 25 and 35 years. Just over one-quarter (27.9%) were co-administered influenza vaccine. A range of pre-existing conditions were present among 18% of participants, including pregnancy-related conditions (2.0%)

including pre-eclamptic toxemia (PET), symphysis pubis dysfunction, low lying placenta, shortening cervix and poor fetal growth. Participant demographics are presented in [table 1](#).

Approximately one-half of Northern participants were co-administered influenza vaccine and 13% of Canterbury participants; this was due to the differences in timing for the influenza season with recruitment.

Injection site reactions

Pain was the most commonly reported reaction to the Tdap injection with 79.0% of participants overall reporting mild or moderate pain in total. Severe pain was reported by 2.6%. Onset of pain occurred within 24 h in 83.9% of participants. There were differences in the intensity, onset and resolution. Northern study participants reported higher intensity pain, later onset and longer time until resolution ([table 2](#)). The proportion reporting no pain was similar for both groups.

Swelling at the injection site was uncommon with 7.6% of participants reporting any swelling and three participants (0.4%) reporting swelling greater than 5 cm. All cases of swelling had an onset within 48 h with half unresolved after 48 h ([table 3](#)). The timing of onset of swelling reported was different between the two study populations with 94% of the Southern population reporting the onset of swelling within 24 h compared with 48% of the Northern population. The time to





Figure 3 Classification of labour, delivery and infant-related adverse events.



*Those who were contacted but ineligible were: 84 women who received Tdap outside the specified inclusion criteria; four women who delivered prior to being contacted; and two males.

Figure 4 Participant flow.

Table 1 Demographic characteristics of participants

	n (%)
Ethnicity (N=793)	
NZ European	583 (73.5)
Māori (indigenous NZ)	55 (6.9)
Pacific Island	34 (4.3)
Asian	103 (13.0)
Middle Eastern, Latin American or African	18 (2.3)
Age (years)	
<20	12 (1.5)
20–24	65 (8.2)
25–29	180 (22.7)
30–34	307 (38.7)
35–39	180 (22.7)
40 or more	49 (6.6)
Co-administered flu vaccine	
Yes	218 (27.5)
Medical history/pre-existing conditions	
None noted	651 (82.1)
Asthma	24 (3.0)
Atopy	4 (0.5)
Cancer	2 (0.3)
Cardiovascular	12 (1.5)
Chronic renal disease	1 (0.1)
Chronic respiratory disease	1 (0.1)
Diabetes	29 (3.7)
Other	52 (6.6)
Pregnancy-related conditions	16 (2.0)
Unanswered	1 (0.1)

NZ, New Zealand.

resolution was the same in both groups with 49% resolving within 48 h (not shown).

Erythema was reported by 5.8% of participants with 0.4% reporting erythema of greater than 5 cm. Onset was generally within 48 h and resolution over 48 h for half of these (table 3). While there were 37 reports of erythema for the Northern participants, there were 9 reports for the Southern participants.

Induration as an injection site reaction was collected in the Northern but not the Canterbury arm. Any induration was reported by 12% of participants, of whom half took more than 48 h to resolve. No induration events measured greater than 5 cm (table 3).

Systemic events

Fever was reported by 17 (2.1%) participants, 14 of whom reported this occurring within 24 h. Of those reporting fever within 24 h, 6 (35%) had been co-administered influenza vaccine (table 4). There were 10 reports of fever in the Northern participants and 7 reports in the Southern participants, and 6/10 Northern participants and 0/7 Southern participants were co-administered influenza vaccine.

Other systemic events included headache and dizziness, nausea and vomiting, fatigue and myalgia or arthralgia. All were uncommon and reported by fewer than 4% of participants with the exception of fatigue, which was reported by 8.4% (table 5). Thirty-two participants reported more than one of these outcomes occurring

Table 2 Number and percentage (n/%) of participants from respective study arms reporting pain after Tdap injection

	Northern arm (N=326)	Canterbury arm (N=467)	Total (N=793)
Pain			
None	56 (17.2)	90 (19.3)	146 (18.4)
Mild, still able to move arm normally	163 (50.0)	184 (39.4)	347 (43.8)
Moderate, hurts to move or to touch	90 (27.6)	189 (40.5)	279 (35.2)
Severe, unable to move arm	17 (5.2)	4 (0.9)	21 (2.6)
Onset of pain (h)	(n=270)	(n=377)	(n=647)
0–24	201 (74.4)	342 (90.7)	543 (83.9)
25–48	68 (25.2)	35 (9.3)	103 (15.9)
49–72	1 (0.4)	–	1 (0.2)
Pain resolved by (h)			
0–24	30 (11.1)	131 (34.7)	161 (25.1)
25–48	86 (31.9)	141 (37.4)	227 (35.4)
>49	154 (57.0)	99 (26.3)	253 (39.5)
Missing	–	6 (1.6)	6 (0.9)

together. Around one-quarter to one-third of those who reported a systemic event also received the influenza vaccine at the same time (table 5).

AEs and SAEs

During the study period, there were 115 AEs and SAEs in 113 participants (two events each for two participants) reported to CARM, in accordance with local guidelines. CARM reports submitted for events not considered as SAEs were mostly minor in nature and for a range of issues including a stiff neck on the same side as the injection, itching, changes in baby movements and combinations of systemic events such as headaches, nausea, fatigue and fever (see tables 4 and 5). Additionally, 25

reports were for diagnosed non-injection site infections (eg, chest, urine and throat).

We have reported SAEs according to whether they occurred or were triggered during pregnancy; during labour and delivery; or occurred in the infant after delivery.

Of the 31 events deemed to be serious (3.9%), there were 23 hospitalisations following immunisation that occurred during pregnancy. Reasons for these were: obstetric bleeding (4), hypertension (2), infection (4), tachycardia (1), preterm labour (9), exacerbation of pre-existing condition (2) and pre-eclampsia (1). All had variable onset time from vaccination (table 6).

Additionally, there were a total of eight SAEs that occurred during labour and delivery: six reported in the Northern arm and two in the Canterbury arm (note the different methodologies). Of these eight SAEs,

two were perinatal deaths, one of which was due to a congenital abnormality, the other unexplained. There was one cyanotic episode and five cases where concern for fetal well-being resulted in health service intervention (table 7).

Table 3 Total number and per cent (n/%) of participants reporting swelling, erythema or induration after Tdap injection

	Swelling (N=793)	Erythema (N=793)	Induration* (N=326)
None	733 (92.4)	747 (94.2)	287 (88.0)
Circumference of event (cm)	(n=60)	(n=46)	(n=39)
>0.0 to <1.0	33 (4.2)	25 (3.2)	21 (6.4)
>1.0 to <2.5	12 (1.5)	10 (1.3)	12 (3.7)
>2.5 to <5.0	12 (1.5)	8 (1.0)	6 (1.8)
>5.0 to <10.0	2 (0.3)	3 (0.4)	–
>10.0 to <15.0	1 (0.1)	–	–
Onset of event after injection (h)	(n=60)	(n=46)	(n=39)
0–24	44 (73.3)	27 (58.7)	15 (38.5)
25–48	16 (26.7)	16 (34.8)	22 (56.4)
>49	–	3 (6.5)	2 (5.1)
Resolution of event (h)	(n=60)	(n=46)	(n=39)
0–24	14 (23.3)	13 (28.3)	6 (15.4)
25–48	15 (25.0)	10 (21.7)	11 (28.2)
>49	31 (51.7)	23 (50.0)	22 (56.4)

*Northern study data only.

Table 4 Numbers and percentage (n/%) of participants reporting fever and taking antipyretics or pain medication following Tdap injection

Fever (N=793)	
No reported fever	776 (97.9)
Fever	17 (2.1)
Onset of fever (n=17) (h)	
0–24	14 (82.3)
25–48	2 (11.8)
49–72	1 (5.9)
Antipyretic or pain medication taken (N=793)	
No	762 (96.1)
Yes	31 (3.9)
Fever within 24 h and co-administered influenza vaccine (n=17)	
Yes	6 (43)

Table 5 Number and percentage of participants reporting systemic events following Tdap injection

Systemic events (N=793)	n (%)	Onset within 24 h n (%)	Co-administered flu vaccine	
			n (%) of sample experiencing event	n (%) of sample with 24 h onset
Headache/dizzy	31 (3.9)	27 (87)	13 (42)	11 (41)
Nausea/vomiting	22 (2.8)	19 (86)	5 (23)	5 (26)
Fatigue	67 (8.4)	58 (86)	17 (25)	13 (22)
Myalgia/arthralgia	24 (3.0)	21 (88)	6 (25)	6 (29)

DISCUSSION

Reduced-antigen-content tetanus–diphtheria–acellular pertussis (Tdap) vaccines have been shown to be highly effective in pregnancy for reducing infant pertussis morbidity,^{3 4} and are being increasingly recommended on national schedules, particularly during epidemics. While there have been no safety concerns to date using these vaccines in pregnancy, data on safety in pregnancy are relatively limited. This study intensively followed 793 women vaccinated in pregnancy with Tdap for solicited and unsolicited outcomes. Our sample included pregnant women regardless of underlying conditions, including women with comorbidities. Injection site reactions were common, minor and self-limiting. Systemic reactions were uncommon. Differences in pain reporting between

the study groups are likely due to both demographic differences and differences in data collection methods. Since there was delay in interviewing the Southern participants and no participant-held diary, minor events are possibly prone to recall bias. Previous NZ studies have shown that vaccine reactogenicity varies by ethnic group.^{20 21} SAEs occurred in our study population during the study period. None were likely to have been caused by the exposure to Tdap vaccine.

Since this is an observational study with no direct comparator groups, the size of the study has limited power to detect SAEs. Data collection methods and periods for each region were different and the Northern region included a wider gestational age band. While our results add to the body of evidence of safety for pregnant

Table 6 SAEs reported among pregnant women following immunisation in 793 pregnant women vaccinated with Tdap

Participant	Event	SAE definition	Onset post-Tdap (days)	Gestation at time of event (weeks)
1	PV bleeding	Required hospitalisation	11	30
2	Hypertension in pregnancy	Required hospitalisation	27	40+
3	Pelvic pain, bacterial vaginitis	Required hospitalisation	9	34
4	Cellulitis	Required hospitalisation	6	38
5	Hypertension in pregnancy	Required hospitalisation	11	34
6	Threatened labour and group A strep infection	Required hospitalisation	20	31
7	Bleeding	Required hospitalisation	24	36
8	Maternal tachycardia	Required hospitalisation	16	34
9	Gestational diabetes and antenatal partum haemorrhage	Required hospitalisation	29	34
10	Preterm labour	Required hospitalisation	1	36
11	Preterm labour	Required hospitalisation	24	36
12	Preterm labour	Required hospitalisation	7	33
13	Exacerbation of pre-existing condition	Required hospitalisation	17	36
14	Exacerbation of pre-existing condition	Required hospitalisation	7	19
15	Pre-eclampsia	Required hospitalisation	8	36
16	Preterm labour	Required hospitalisation	16	36
17	Preterm labour	Required hospitalisation	19	36
18	Preterm labour	Required hospitalisation	11	33
19	Preterm labour	Required hospitalisation	19	36
20	Vaginal bleeding	Required hospitalisation	21	34
21	Preterm labour	Required hospitalisation	9	34
22	Preterm labour	Required hospitalisation	~15*	36
23	Infection (coronavirus)	Required hospitalisation	1	34

*No delivery date was recorded for this participant.
SAE, serious adverse event.

Table 7 SAEs during labour in and among infants of 793 pregnant women vaccinated with Tdap

Participant	Event	SAE definition	Onset post-Tdap (days)	Gestation at time of event (weeks)
24	Cyanotic episodes in infant	Prolongation of existing hospitalisation	31	39
25	Fetal death (trisomy 11q)	Resulted in death	10	36
26	Non-reassuring fetal status	Intervention to prevent permanent damage or death	11	38
27	Non-reassuring fetal status	Hospitalisation	23	39
28	Non-reassuring fetal status	Intervention to prevent permanent damage or death	37	41
29	Non-reassuring fetal status	Intervention to prevent permanent damage or death	18	39
30	Non-reassuring fetal status	Admission to the NICU and continuous positive airway pressure	28	37
31	Perinatal death (stillbirth)	Resulted in death	53	40

Clinician review and assessment of each of these cases found that none were likely to be vaccine related. NICU, neonatal intensive care unit; SAE, serious adverse event.

women, the study population was not randomly selected. The women in our group were older (32 vs 29.2 years) and more likely to be of NZ European ethnicity (73.5% vs 49.5%) than the general NZ maternal population. This is most likely a reflection of the health-seeking behaviour associated with these demographics. While pregnancy complications increase with age, in contrast, NZ European ethnicity is associated with lower risk pregnancies.²²

The reactogenicity of acellular pertussis-containing vaccines in adults was reviewed in 2012.²³ Data on the reactogenicity of Tdap are derived from clinical trials and prospective studies; however, the results are not presented in a standardised way that allows for comparison. There is insufficient information in the study manuscripts to determine in more detail how events are measured or classified. However, severe local pain ranges from less than 1% to 30%, and severe local redness or swelling (≥ 50 mm) ranges from 2% to 18%.²⁴⁻³⁰ Although most of these studies were published after 2007 when definitions were published by the Brighton Collaboration, none have provided information about induration.

A small US randomised trial including 33 pregnant women assessed the immunogenicity and safety of Tdap during pregnancy. Safety outcomes were collected with a 7-day diary. How these were defined is not described. Pain was reported by 25 (75.8%) participants while erythema and induration/swelling were each reported by 3 (9.1%) participants. These were the same rates as reported by the non-pregnant women in the study.³¹ The rates of injection site pain in our study are consistent with most other studies that report less than 10% experiencing severe pain.^{16 25 26 28} Rates of swelling with a diameter of more than 50 mm ranged from 1% to 18% in clinical studies. In our study population, just 0.4% recorded a swelling of more than 50 mm and

fewer than 8% reported any.^{16 24-28 30} Our outcomes are lower than these other reports; however, this could be because we have differentiated between swelling and induration. Overall, both events were infrequent and mild. Erythema (redness) was relatively uncommon in our study with fewer than 6% of participants reporting any and 1.4% reporting a diameter greater than 50 mm. In clinical studies, severe erythema has been reported to range from 2% to 17%.^{24-28 30} Our rates are consistent with those recently reported in the US trial in pregnant women.³¹ It should be noted that the aluminium content of the NZ Boostrix formulation is higher than that of the US formulation, and the higher aluminium content may be linked to greater local reactogenicity.³²

In the Northern arm of our study, we collected information about swelling and induration according to Brighton Collaboration definitions^{17 18} and are able to report induration separately from swelling. Induration occurred more frequently than swelling and appeared to have a later temporal onset than swelling. More than half of the cases of induration occurred 25–48 h later compared with around a quarter of all swellings, supporting the likelihood that each has a different aetiology. It is likely that some of the swelling reported in the Southern participants was in fact induration misclassified.

We found differences in the reporting of swelling and erythema between our groups. There were fewer reports of erythema in the Southern group, and while there were similar numbers of reports of swelling, the timing for onset may have been more imprecise. Given the differences in the reported pain onset and resolution between the groups, it is likely that recall for details of minor local reactions became more prone to recall bias as time progressed and in the absence of a diary to record the details.

While in previous clinical studies headache has tended to be reported by around a third of participants,^{24 25 27–30} fewer than 3.9% of our participants reported either headache or feeling dizzy. Also, few of our participants reported any gastrointestinal symptoms (2.8%). Almost all systemic events in our study occurred in the 24 h following immunisation as opposed to later, and we consider them likely to be vaccine related. Overall, there were few systemic events reported and the rates of the most common are consistent with those reported in the US study with 33 participants.³¹ In our study, one-third to one-quarter of those who reported systemic events had also received the influenza vaccine at the same time. Northern participants reporting systemic events were consistently more likely to have received co-administered influenza vaccine than Southern participants. Most of the reported fevers in our Northern participants also occurred after co-administration with influenza vaccine. Some influenza vaccines are known to be more pyrogenic than others, so it is possible that influenza vaccine was the cause of these excess reports, particularly as the fevers occurred within 24 h after vaccination.²¹ The safety of co-administration of influenza vaccine and Tdap in pregnancy has been assessed by the Vaccine Safety Datalink Project, and no excess medically attended events occurred among women receiving both vaccines together compared with sequential vaccination.⁶ However, it cannot be assumed that these vaccines have the same reactogenicity profiles.

SAEs occurred in our study population during the study period. None were considered by clinical review as likely to be caused by the exposure to Tdap vaccine. In NZ, up to 15% of pregnancies have obstetric complications, and approximately 1 per 10 infants are born preterm or low birth weight.^{33 34} Annually, there are approximately 600 perinatal deaths (~1% of births) of which 14% are unexplained. These figures have remained consistent over time.²² On the basis of NZ data, the rates of SAEs in our study were not higher than the expected background rate for such a cohort. Reports to the US Vaccine Adverse Event Reporting System of pregnant women inadvertently given Tdap have been summarised.³⁵ Between January 2005 and June 2010, there were 132 reports identified, 20 following Boostrix with no non-pregnancy SAEs reported. In the US trial,³¹ SAEs occurred in 7/33 pregnant women, without known risks to pregnancy at enrolment, followed to 4 months postpartum, and none of these were non-pregnancy SAEs. The events occurred at variable time periods following immunisation and none were considered attributable to the vaccine. Recently, obstetric events and birth outcomes for 123 494 women from two Californian Vaccine Safety Datalink sites were evaluated; 26 229 women received Tdap with no increased risk for hypertensive disorders of pregnancy or preterm or small for gestational age birth found. There was a small increased risk of chorioamnionitis diagnosis.⁹ Further investigations have not found an association.^{7 8} A matched

cohort study from the UK in 20 074 pregnant women and a matched historical unvaccinated control group found no evidence of any increase for predefined pregnancy-related AEs including stillbirth.¹⁰ These studies all support the safety profile of Tdap in pregnancy.

In conclusion, we found that a Tdap vaccination was well tolerated in pregnant women. Our findings are consistent with data from studies involving non-pregnant women.²³ There were no SAEs in this study that were likely to have been caused by the vaccine. This is reassuring for pregnant women, vaccinators and policymakers.

Twitter Follow Helen Petousis-Harris at @PetousisH and Janine Paynter at @JaninePaynter

Acknowledgements The authors acknowledge the general practice and DHB antenatal clinic staff involved in the study, for their assistance in referring eligible women to the study team. They thank the pregnant women and mothers who contributed to the study. They also thank their colleagues Angela Chong, Trish Graham, Nicky Ellis, Barbara McArdle, Tracey Poole and Jane Stephen who assisted in various ways, such as recruiting general practices, database set-up and data collection.

Contributors HP-H and TW were responsible for the study concept and design. DW acquired the data for the Northern arm of the study. PG acquired the data for the Canterbury arm of the study. JP managed and analysed the data for this study. NT collected clinical data for the SAEs. All authors drafted the manuscript and critically revised it for important intellectual content.

Funding The Northern study was funded by GlaxoSmithKline Biologicals SA, the manufacturer of Boostrix and sponsored by Auckland UniServices Ltd. The Southern study was funded by the Canterbury District Health Board.

Competing interests All authors have completed the ICMJE uniform disclosure form at http://www.icmje.org/coi_disclosure.pdf and declare: financial support for the submitted work from GlaxoSmithKline. HP-H has served on advisory panels and/or DSMB for GlaxoSmithKline, Pfizer, CSL Biotherapies and Merck. She has also served on investigator-led studies funded by GlaxoSmithKline, CSL and sanofi pasteur.

Ethics approval Ethics approval was obtained from the University of Auckland Human Participants Ethics Committee (010656) and from the Upper South A Regional Ethics Committee (URA/12/EXP/021).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Advisory Committee on Immunization Practices (ACIP). *Summary Report February 23–24, 2011*. Atlanta, Georgia, 2011.
2. Davies SC. Temporary programme of pertussis (whooping cough) vaccination of pregnant women. 28 Sept 2012. Department of Health, London, 2012.
3. Amirhalingam G, Andrews N, Campbell H, *et al*. Effectiveness of maternal pertussis vaccination in England: an observational study. *Lancet* 2014;384:1521–8.
4. Dabrera G, Amirhalingam G, Andrews N, *et al*. A case-control study to estimate the effectiveness of maternal pertussis vaccination in protecting newborn infants in England and Wales, 2012–2013. *Clin Infect Dis* 2015;60:333–7.
5. Englund JA. Maternal immunization—promises and concerns. *Vaccine* 2015;33:6372–3.

6. Sukumaran L, McCarthy NL, Kharbada EO, *et al.* Safety of tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis and influenza vaccinations in pregnancy. *Obstet Gynecol* 2015;126:1069–74.
7. Morgan JL, Baggari SR, McIntire DD, *et al.* Pregnancy outcomes after antepartum tetanus, diphtheria, and acellular pertussis vaccination. *Obstet Gynecol* 2015;125:1433–8.
8. Datwani H, Moro PL, Harrington T, *et al.* Chorioamnionitis following vaccination in the Vaccine Adverse Event Reporting System. *Vaccine* 2015;33:3110–13.
9. Kharbada EO, Vazquez-Benitez G, Lipkind HS, *et al.* Evaluation of the association of maternal pertussis vaccination with obstetric events and birth outcomes. *JAMA* 2014;312:1897–904.
10. Donegan K, King B, Bryan P. Safety of pertussis vaccination in pregnant women in UK: observational study. *BMJ* 2014;349:g4219.
11. Moro PL, McNeil MM, Sukumaran L, *et al.* The Centers for Disease Control and Prevention's public health response to monitoring Tdap safety in pregnant women in the United States. *Hum Vaccin Immunother* 2015;11:2872–9.
12. Ministry of Health. Processes for safe immunisation. In: Ministry of Health, ed. *Immunisation Handbook 2014*. Wellington: Ministry of Health, 2014:75.
13. International Conference on Harmonisation. Clinical safety data management: definitions and standards for expedited reporting E2A. *International Conference on Harmonisation*; 1994.
14. International Conference on Harmonisation. Post-approval safety data management: definitions and standards for expedited reporting E2D. *International Conference on Harmonisation*; 2003.
15. Working Group on Vaccine Pharmacovigilance. *Definition and application of terms for vaccine pharmacovigilance*. Geneva, Switzerland: World Health Organization, 2012.
16. Gidudu J, Kohl KS, Halperin S, *et al.* A local reaction at or near injection site: case definition and guidelines for collection, analysis, and presentation of immunization safety data. *Vaccine* 2008;26:6800–13.
17. Kohl KS, Walop W, Gidudu J, *et al.* Swelling at or near injection site: case definition and guidelines for collection, analysis and presentation of immunization safety data. *Vaccine* 2007;25:5858–74.
18. Kohl KS, Walop W, Gidudu J, *et al.* Induration at or near injection site: case definition and guidelines for collection, analysis, and presentation of immunization safety data. *Vaccine* 2007;25:5839–57.
19. Michael Marcy S, Kohl KS, Dagan R, *et al.* Fever as an adverse event following immunization: case definition and guidelines of data collection, analysis, and presentation. *Vaccine* 2004;22:551–6.
20. Petousis-Harris H, Jackson C, Stewart J, *et al.* Factors associated with reported pain on injection and reactogenicity to an OMV meningococcal B vaccine in children and adolescents. *Hum Vaccin Immunother* 2015;11:1875–80.
21. Petousis-Harris H, Poole T, Turner N, *et al.* Febrile events including convulsions following the administration of four brands of 2010 and 2011 inactivated seasonal influenza vaccine in NZ infants and children: the importance of routine active safety surveillance. *Vaccine* 2012;30:4945–52.
22. Perinatal and Maternal Mortality Review Committee. *Eighth annual report of the perinatal and maternal mortality review committee*. Reporting mortality 2012. Wellington, 2014.
23. McCormack PL. Reduced-antigen, combined diphtheria, tetanus and acellular pertussis vaccine, adsorbed (Boostrix®): a review of its properties and use as a single-dose booster immunization. *Drugs* 2012;72:1765–91.
24. Abarca K, Valdivieso F, Potin M, *et al.* [Immunogenicity and reactogenicity of a reduced antigen content diphtheria, tetanus and acellular pertussis vaccine (dTpa) in 10 to 11 years old children and in adults]. *Rev Med Chil* 2002;130:502–10.
25. Blatter M, Friedland LR, Weston WM, *et al.* Immunogenicity and safety of a tetanus toxoid, reduced diphtheria toxoid and three-component acellular pertussis vaccine in adults 19–64 years of age. *Vaccine* 2009;27:765–72.
26. Booy R, Van Der Meeren O, Ng SP, *et al.* A decennial booster dose of reduced antigen content diphtheria, tetanus, acellular pertussis vaccine (Boostrix™) is immunogenic and well tolerated in adults. *Vaccine* 2010;29:45–50.
27. Chan S, Tan PT, Han HH, *et al.* Immunogenicity and reactogenicity of a reduced-antigen-content diphtheria-tetanus-acellular pertussis vaccine as a single-dose booster in Singaporean adults. *Singapore Med J* 2006;47:286–90.
28. Mertsola J, Van Der Meeren O, He Q, *et al.* Decennial administration of a reduced antigen content diphtheria and tetanus toxoids and acellular pertussis vaccine in young adults. *Clin Infect Dis* 2010;51:656–62.
29. Tumbull FM, Heath TC, Jalaludin BB, *et al.* A randomized trial of two acellular pertussis vaccines (dTpa and pa) and a licensed diphtheria-tetanus vaccine (Td) in adults. *Vaccine* 2000;19:628–36.
30. Van der Wielen M, Van Damme P, Joossens E, *et al.* A randomised controlled trial with a diphtheria–tetanus–acellular pertussis (dTpa) vaccine in adults. *Vaccine* 2000;18:2075–82.
31. Munoz FM, Bond NH, Maccato M, *et al.* Safety and immunogenicity of tetanus diphtheria and acellular pertussis (tdap) immunization during pregnancy in mothers and infants: a randomized clinical trial. *JAMA* 2014;311:1760.
32. Lindblad EB. Aluminium adjuvants—in retrospect and prospect. *Vaccine* 2004;22:3658–68.
33. Ministry of Health. *New Zealand Maternity Clinical Indicators 2013*. Wellington: Ministry of Health, 2015.
34. Ministry of Health. *Hospital-based Maternity Events 2007*. Wellington: Ministry of Health, 2010.
35. Zheteyeva YA, Moro PL, Tepper NK, *et al.* Adverse event reports after tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis vaccines in pregnant women. *Am J Obstet Gynecol* 2012;207:59.e1–7.